

UCCE Delta Sorghum Seeding Rate Trial – 2016-17 Final Report

Michelle Leinfelder-Miles, Delta Farm Advisor

In recent years, the University of California Division of Agriculture and Natural Resources has augmented research efforts on growing grain and silage sorghum [*Sorghum bicolor* (L.) Moench] in California. The purpose of the Delta Sorghum Seeding Rate Trial was to better understand optimal seeding rates for grain sorghum grown in the Sacramento-San Joaquin River Delta. While such information exists for Midwest sorghum production¹, applied information is lacking for California, and more specifically for the Delta – a unique agricultural region known for its organic soils, shallow groundwater, and cooler climate conditions. This information is important because sorghum has similar growth habits as corn and is sometimes grown as a substitute for corn because of its tolerance of drought and low-input conditions. In the United States, sorghum is used in a wide array of feedstocks for biofuels, pet foods, dairy, cattle, pork and poultry feed, and more recently as a gluten-free cereal grain for human food systems.

The trial took place during the 2016 and 2017 growing seasons on Tyler Island in Sacramento County. In 2016, the plot was established on a Rindge mucky silt loam with approximately 20 percent organic matter in the top 15 inches of soil. In 2017, the plot was established on a Gazwell mucky clay with approximately 10 percent organic matter in the top 30 inches of soil. Approximately 55,600 acres in the Delta are described by the Rindge classification, and approximately 24,600 acres are described by the Gazwell classification. The 2016 trial was planted on May 20th, and the 2017 trial was planted on May 25th using a John Deere cone planter. Seed was planted approximately 2 inches deep. We used the grower's varieties, which were Eureka Seeds 3292 in 2016 and Eureka Seeds 3325W in 2017. Both varieties were white sorghum varieties, had 16,000 seeds/lb, and 85 percent germination, according to the labels. Five seeding rate treatments (5, 6, 9, 12, and 15 lbs/acre) were tested. In 2016, the five treatments were replicated over four blocks positioned down the rows in a Randomized Complete Blocks Design. In 2017, the five treatments were replicated five times in a Latin Square design. The experimental design was changed to a Latin Square in 2017 in order to control field variability both down and across the rows. Each plot consisted of four rows (30-inch row spacing) that were 45 feet in length in 2016 and 50 feet in length in 2017. In 2016, the previous crop in the field was wheat, and in 2017, the previous crop was corn. The fields were managed similarly in both years. Subsurface irrigation by "spud ditch" was employed twice; however, the amount of applied water is difficult to ascertain with this method of irrigation. The fertility program was 35 gallons/acre of 8-24-0 with ½ percent of zinc at planting. The field was cultivated one time, and Maestro 4EC (8 oz/acre), AAtrex 4L (0.75 pint/acre), and Crosshair (4 oz/acre) were applied for post-emergence weed control in mid-June. The plots were harvested on November 14, 2016 and October 12, 2017 using an Almaco research combine, harvesting the center two rows from the four-row plots.

Trial results are presented as plant establishment characteristics (Table 1), plant maturity characteristics (Tables 2 and 3), and yield (Figures 1). The tables and figure present mean values for the four (2016) or five (2017) replicates. Tukey's range test was the statistical method used to compare the means. Treatments were considered statistically different if their P value was less than 0.05, or 5 percent. Differences among treatments are indicated by different letters following the mean.

The seeding rates are expressed as plant populations in Table 1. The number of sorghum seeds/lb is highly variable across varieties. For this reason, when determining seeding rates, growers should first determine their

¹ Examples from Wisconsin (<http://corn.agronomy.wisc.edu/Crops/SorghumGrain.aspx>), Alabama (<http://www.aces.edu/pubs/docs/A/ANR-0502/index2.tmpl>), and Texas (<http://sanangelo.tamu.edu/extension/agronomy/agronomy-publications/grain-sorghum-production-in-west-central-texas/#planting>).

desired plant population. A worksheet at the end of this report provides equations for calculating seeding rate based on desired plant populations and percent germination for the variety. Stand counts were made as the number of plants/10-foot row length approximately two weeks and one month after planting. The counts were scaled up to plants/acre. Across both years, stands generally decreased from the first count date to the second. Stand counts were lower in 2017 compared to 2016, but this did not translate into lower yields. Weeds were also counted in the month after planting (data not shown), but overall weed pressure was very low in both years.

Table 1. Plant establishment characteristics of the 2016 and 2017 UCCE Delta sorghum seeding rate trial.

Seeding Rate (lbs/acre)	Estimated Plant Population (# seeds/acre)	2016 Stand Counts		2017 Stand Counts	
		1-June (plants/acre)	16-June (plants/acre)	13-June (plants/acre)	30-June (plants/acre)
5	80,000	82,756 e	79,489 e	69,515 d	68,818 e
6	96,000	106,712 d	96,258 d	76,832 d	86,763 d
9	144,000	149,396 c	130,667 c	124,395 c	111,154 c
12	192,000	196,436 b	161,156 b	161,679 b	151,399 b
15	240,000	248,267 a	190,338 a	200,705 a	187,463 a
Treatment P value		<0.0001	<0.0001	<0.0001	<0.0001
Standard Error		4432	3748	4812	5760

There were no differences in the number of days to flowering among treatments in either year; however, there were differences among treatments in the other maturity characteristics. In both years, plants in the higher seeding rate plots were taller and/or had longer panicle exertion (the length of the stem from the top leaf to the bottom of the panicle), suggesting that at higher densities, plants were competing with each other and growing longer internodes. In both years, panicles were longest in the 5 lb seeding rate and statistically longer than the panicles in the higher rate treatments. There were no statistical differences in grain moisture at harvest in either year. In 2017, the weight of 1000 seeds was also analyzed to understand whether certain treatments were producing heavier seeds. We found that the seed of the 6 lb rate had the heaviest weight and was statistically heavier than the 12 lb rate but not different from the other treatments.

Table 2. Plant maturity characteristics of the 2016 UCCE sorghum seeding rate trial in the Delta.

Seeding Rate (lbs/acre)	Days to Flowering (# of days)	Plant Height (inches)	Panicle Length (inches)	Panicle Exsertion (inches)	Moisture (%)
5	71	53.1 b	11.8 a	6.2 c	18.2
6	71	52.3 b	11.0 abc	6.7 c	18.0
9	71	54.7 a	11.1 ab	7.4 bc	17.2
12	71	53.4 ab	10.6 bc	8.1 ab	17.8
15	71	53.7 ab	10.3 c	9.1 a	18.1
Treatment P value	0.9014	0.0003	<0.0001	<0.0001	0.0589
Standard Error	0.4	0.8	0.3	0.4	0.22


Table 3. Plant maturity characteristics of the 2017 UCCE sorghum seeding rate trial in the Delta.

Seeding Rate (lbs/acre)	Days to Flowering (# of days)	Plant Height (inches)	Panicle Length (inches)	Panicle Exsertion (inches)	Moisture (%)	1000 seed weight (grams)
5	70	59 ab	10.9 a	6.8 c	12.1	19.3 ab
6	70	60 ab	10.8 ab	7.3 bc	12.4	19.4 a
9	70	60 ab	10.2 bc	8.0 ab	12.0	19.3 ab
12	70	61 a	10.0 c	8.6 ab	13.0	18.9 b
15	70	59 b	10.2 bc	8.4 a	12.2	19.0 ab
Trt. P value	0.0685	0.0264	<0.0001	<0.0001	0.1749	0.0314
Standard Error	0.57	0.54	0.45	0.55	0.29	0.19

While there were no statistical differences in yield across treatments in either year, the take-home message of the trial is that there appears to be no benefit to planting the highest seeding rates. In both years, the trend was for the 15-lb seeding rate to have the lowest yield. In 2016, there was a lot of variability in the data. There was a trend for the 9-lb treatment to have higher yields; however, we suspect this was due to the experimental design. In 2016, by random chance, there were several 9-lb treatment plots next to the sub-irrigation ditches, which were exterior to the experiment on both sides. For this reason, the 9-lb treatment may have been inadvertently favored with better moisture conditions. To correct for this, the experimental design was improved in 2017 by blocking in both directions, in order to better control field variability. The 2017 yields were consistent across treatments, around 7000 lbs/acre. The 2017 results best illustrate how planting the higher seeding rates provided no yield benefit, yet incurs a higher seed expense. We recognize that growers will need to consider site characteristics, like weed or wireworm pest pressure, when determining optimal seeding rates; nevertheless, this research indicates that good

yields can result from seeding rates of 5 or 6 lbs/acre (estimated plant populations of 80-96,000 plants/acre), and that planting higher plant populations would not only cost growers more in seed expense but could also cost them in yield.

Figure 1. Yield at 13 percent moisture of UCCE Delta sorghum seeding rate trial. There were no significant differences among treatments in 2016 ($P = 0.1278$) or 2017 ($P = 0.2419$).



In summary, it is important to study sorghum cultural practices in the Sacramento-San Joaquin Delta region, and in California at-large, because currently, most applied information comes from the Midwest. California growers need information on sorghum cultivation because sorghum may be grown as a lower-input substitute for corn. Sorghum seeding rates were studied to assist growers with determining optimum rates for the Delta environment. The results indicate that there is no yield benefit to planting seeding rates greater than 6 lbs/acre (estimated plant population greater than 96,000 plants/acre), and that planting higher rates is just added expense for the grower. Future research should investigate these plant populations on narrower row spacing. Special thanks go to growers, Steve and Gary Mello, and to UC Kearney Research and Extension Center Director, Jeff Dahlberg, for providing equipment and information for the success the trial.

Sorghum Plant Population Worksheet

Jeff Dahlberg, Director, Kearney Agricultural Research and Extension Center

Calculating number of seeds per foot to get desired plant population per acre:

Table 1. Sorghum plant population in plants/ft-row for different row spacings.

	Desired plants per acre (X1000)					
	25	50	75	100	125	150
Row width	Average number of plants per foot of row					
7 inch	0.3	0.7	1.0	1.3	1.7	2.0
15 inch	0.7	1.4	2.2	2.9	3.6	4.3
20 inch	1.0	1.9	2.9	3.8	4.8	5.7
30 inch	1.4	2.9	4.3	5.7	7.2	8.6
40 inch	1.9	3.8	5.7	7.7	9.6	11.5

How do you calculate plant population and seeding rate?

Plant population per acre is equal to:

$$[43,560 \div (\text{row spacing in inches} \div 12)] \times \text{plants per foot of row}$$

Seeding rate per acre is equal to:

$$\frac{\text{Suggested plants per foot of row}}{\text{Percent germination} \times \text{Percent expected emergence}}$$